

Bidicod

Butamirate Citrate BP

Composition

Bidicod Syrup: Each 5 ml syrup contains Butamirate Citrate BP 7.5 mg.

Pharmacology

Butamirate Citrate exerts expectorant, moderate bronchodilating and anti-inflammatory action. It is safe and non-sedating which is neither chemically nor pharmacologically related to opium alkaloids. The active metabolites of Butamirate Citrate have also antitussive action.

Indication

Bidicod is indicated in acute cough of any etiology, suppression of pre- & post-operative cough in surgery & bronchoscopy and suppression of whooping cough.

Dose and administration

Route of administration: Oral

Bidicod Syrup: Children 3-6 years of age: 5 ml 3 times daily.

Children 6-12 years of age: 10 ml 3 times daily.

Adolescent (over 12 years of age): 15 ml 3 times daily.

Adults: 15 ml 4 times daily.

Contraindication

Butamirate Citrate is contraindicated in patients with known hypersensitivity to any component of the formulation.

Warning & Precaution

Avoid simultaneous administration of expectorants. Avoid during pregnancy (1st 3 months).

Side effects

Common: Rash.

Rare: Nausea, diarrhoea and vertigo have been observed in a few rare cases, resolving after dose reduction or treatment withdrawal.

Use in pregnancy and lactation

Butamirate Citrate should not be used during the first trimester of pregnancy. During the remainder of pregnancy, it can be used if indicated by a physician but with caution. As a general rule, for safety reasons, in the absence of data on elimination of the active substance in breast milk, the benefits of Butamirate Citrate administration during lactation should be carefully weighed against the risks.

Use in Children & Adolescents

See the dose and administration.

Drug Interaction

With medicine: Concomitant use of expectorants may lead to retention of mucous in the respiratory system, which can increase the risk of bronchospasm and airway infection.

With food & other: No adequate data is found.

Overdose

Accidental overdose with Butamirate Citrate can cause the following symptoms: drowsiness, nausea, vomiting, diarrhoea, loss of balance and hypotension. Standard emergency procedures should be followed: activated charcoal, saline laxatives and standard cardio-respiratory resuscitation.

Storage

Store below 30° C temperature and cool and dry place, away from light. Keep out of the reach of children.

Packing

Bidicod Syrup: Each bottle contains 100 ml Syrup with a measuring cup.



Manufactured by

ONE PHARMA LTD.

C-23-24, BSCIC I/A

Bogura, Bangladesh

LF0049

Rev.02 BDC