

Caftyl

Paracetamol BP & Caffeine BP Tablet

Composition

Caftyl Tablet: Each tablet contains Paracetamol BP 500 mg & Caffeine BP 65 mg.

Pharmacology

Caftyl (Paracetamol & Caffeine) is a fast acting and safe analgesic with marked antipyretic property. It is specially suitable for patients who, for any reason, can not tolerate aspirin or other analgesics. The presence of Caffeine increases the effectiveness of Paracetamol.

Indication

The indications of Caftyl are as follows

- Headache
- Toothache
- Feverishness
- Sore throat
- Helps to reduce temperature
- Migraine
- Neuralgia
- Period pain
- Rheumatic pain & backache
- Pain of colds and flu

Dose and Administration

a) *Route of administration:* Oral

Adult: 1-2 tablets every 4-6 hours. Maximum dose 4 gm (8 tablets) daily.

Contraindication

Paracetamol is contraindicated in patients with severe renal function impairment and hepatic disease (Viral Hepatitis). It is contraindicated in patients with known hypersensitivity to paracetamol or caffeine.

Warning & Precaution

Paracetamol & Caffeine should be given cautiously in the following cases : In patients with hepatic or renal failure, in patients taking other hepatotoxic medication. Prolonged use of the drug without consulting with a physician should be avoided.

Side effects

a) *Common:* Side effects of paracetamol are usually mild, though haematological reactions including thrombocytopenia, leukopenia, pancytopenia, neutropenia and agranulocytosis have been reported.

b) *Rare:* Pancreatitis, skin rashes and other allergic reactions occur occasionally.

Use in pregnancy and lactation

Pregnant mothers should consult with doctors before taking Paracetamol & Caffeine. Paracetamol & Caffeine can be taken while breast feeding.

Use in Children & Adolescents

Not recommended for children under 12 years.

Drug Interaction

a) *With medicine:* Paracetamol & Caffeine increases the effect of chloramphenicol and coumarin anti-coagulant.

b) *With food & others:* Risk of hepatotoxicity of paracetamol may be increased in alcoholics or in patients taking other anti-epileptic medications.

Overdosage

Symptoms of Paracetamol overdose in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 40 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur.

Storage

Store below 30°C. Protect from light and moisture. Keep all medicines out of reach of children.

Packing

Caftyl Tablet: Each box contains 100 (10x10's) tablets in Alu-PVC blister pack.



Manufactured by

ONE PHARMA LTD.

C-23-24, BSCIC I/A

Bogura, Bangladesh

LF0002

CFT Rev.02