

Neobastin 10

Ebastine BP

Composition

Neobastin 10 Tablet: Each film coated tablet contains Ebastine BP 10 mg.

Description

Neobastin 10 (Ebastine) is a long-acting and selective H₁-histamine receptor antagonist. Ebastine is almost totally converted to the pharmacologically active acid metabolite, Carebastine. Peak plasma levels of Carebastine occur at 2.6 to 4 hours and achieve levels of 80 to 100 ng/ml. The half-life of Carebastine is between 15 to 19 hours.

Indications

Ebastine is indicated for the symptomatic treatment of:

- Seasonal and Perennial Allergic Rhinitis
- Chronic Idiopathic Urticaria

Dosage & Administration

Tablet: Adults (more than 12 years of age): 10 mg (one tablet) once daily.

Children (6-12 years of age): 5 mg (half tablet) once daily.

Ebastine may be taken with or without food.

Side Effects

The most common side-effects are headache, dry mouth and drowsiness. Less commonly reported side effects include abdominal pain, dyspepsia, nausea and insomnia.

Precautions

Caution is advised when used in hepatic impairment, renal insufficiency, QTc interval prolongation.

Use in Pregnancy & Lactation

The safety of Ebastine during pregnancy and lactation has not been established.

Drug Interaction

Concomitant use of Ketoconazole, Itraconazole, Clarithromycin or Erythromycin may increase plasma levels of Ebastine and cause QTc interval prolongation.

Overdose

In studies conducted at a high dosage, no clinically meaningful signs or symptoms were observed up to 100 mg given once daily. There is not specific antidote for Ebastine. In case of accidental overdose, gastric lavage, monitoring of vital functions including ECG, and symptomatic treatment should be carried out.

Contraindications

Ebastine is contraindicated to patients with hypersensitivity to any component of this product.

Storage Condition

Store below 30° C temperature. Keep in a cool and dry place, away from light. Keep out of the reach of children.

Commercial Pack

Neobastin 10 Tablet: Each box contains 3 blister of 10 tablets.



Manufactured by

ONE PHARMA LTD.

C-23-24, BSCIC I/A

Bogura, Bangladesh

LF0048

NBN Rev.01